


## Series : 09



42.  $\Delta G$  is quantitative hence it is extensive whereas  $E^\circ_{\text{cell}}$  or  $E_{\text{cell}}$  is intensive property.  $\dots \text{Ans: C}$

43. For zero order reaction  $t = \frac{[\text{R}]_0 - [\text{R}]_t}{k}$   
 $\text{at } t = \frac{1}{2} = \frac{[\text{R}]_0}{2k}$   $t_{100\%} = \frac{[\text{R}]_0}{k}$   
 $\therefore t_{100\%} = 2 \times t_{\frac{1}{2}} = 2 \times 10 = 20 \text{ min} \dots \text{Ans: D}$

44. For 1st order reaction,  $t_{\frac{1}{2}} = \frac{0.693}{k} \Rightarrow k = \frac{0.693}{t_{\frac{1}{2}}}$


$t = \frac{2.303}{k} \log \frac{[\text{R}]_0}{[\text{R}]_t}$  Here  $[\text{R}]_0 = 100$  &  $[\text{R}]_t = 50$

$\therefore t = \frac{2.303}{\frac{0.693}{t_{\frac{1}{2}}}} \log \frac{100}{50}$

$t = \frac{5730}{0.3} \log \frac{100}{50} \therefore \frac{0.693}{2.303} \approx 0.3$   
 $\dots \text{Ans: (A)}$

45. Conceptual  $\dots \text{Ans: A}$

46. For  $n_A \cdot A \rightarrow n_B \cdot B$  Rate =  $-\frac{1}{n_A} \frac{d[A]}{dt} = +\frac{1}{n_B} \frac{d[B]}{dt}$



47. Actual reason is due to smaller size F repels and unlike "O", F forms single bond with Mn. Hence, correct answer  $\dots \text{Option (A)}$

48.  $M^{2+}$  ion has same d electrons value of z on the position of ONE  $\therefore {}_{25}\text{Mn}^{2+} : [\text{Ar}] 3d^5 4s^0$   $\therefore n=5 \Rightarrow \mu_{\text{so}} = \sqrt{35}$   
 $\dots \text{Option (A)}$

49. In Acidic med: Tx BK Example: 9



$\therefore$  Option: (D)

50. Ti has several ox<sup>ion</sup> state in Ziegler-Natta catalyst  
Among these III is the most stable.

$\therefore$  Option: (D)

51. Complex with formula  $[\text{Ma}_3\text{b}_3]$  are most suitable to show facial-meridional isomers.

$\therefore$  In  $[\text{Co}(\text{NH}_3)_x(\text{NO}_2)_y]$ ,  $x$  &  $y$  should be 3

$\therefore x=3, y=3$


$\therefore$  Option: (B)

52.  $[\text{NiCl}_4]^{2-}$  undergo  $sp^3$  hy<sup>l</sup>n &


$[\text{Ni}(\text{CN})_4]^{2-}$  undergo  $dsp^2$  hy<sup>l</sup>n.  $\therefore$  Option: (A)

53. each  $\text{C}_2\text{O}_4^{2-} = -2 \therefore$  trioxalato  $= -6$   $[\text{Al}(\text{OX})_3]^{3-}$   
Al has +3 ox<sup>ion</sup> state

$\therefore$  Formula is  $\text{K}_3[\text{Al}(\text{OX})_3]$  which produces



54. strong ligand  $\propto$  CFSE  $\propto$  E<sub>absorbed</sub>  $\propto \frac{1}{\lambda_{\text{absorbed}}}$



55. Allylic halides have common formula  $\text{C}=\text{C}-\text{C}-\text{X}$

$\therefore$  1-bromobut-2-ene  $\text{Br}-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_3$  is

allylic

$\therefore$  Option (A)

56. mono alkane with optical isomerism  
required chiral carbon i.e. all four different  
valencies.



$\therefore$  Four "C" required.

$\therefore$  Option: (A)




57: Rate of  $S_N2$  is highest in  $1^\circ$  R-X with minimum branching  $\therefore$  1-bromobutane  $\therefore$  Option: (B)

58. Chloroform ( $CHCl_3$ ) Trichloromethane  $\therefore$  Option: (D)

59.



The product has structure  $\therefore$   $R' = CH_3$



$R'' = CH_2CH_3$

$R''' = CH_3$

As this is the one option with total 4 " $^\circ$ "

Other options have 3, 5 total " $^\circ$ "  $\therefore$  Option (A)

60.




61. Glucose  $\xrightarrow{\text{Zymase}}$  ethanol +  $CO_2$   $\therefore$  Option (B)

62. Ethanol  $\xrightarrow[H^+]{[Acid]}$  Diethylether is  $S_N2$

$\therefore$  Option (D)

63. Vanillin is



$\therefore$  It has - phenyl

-  $OCH_3$

-  $CHO$

-  $OH$

$\therefore$  Option (A)

64. PCC is better oxidizing agent to convert  $-CH_2OH$  into  $-CHO$

$\therefore$   $C_5H_5NH^+CrO_3Cl^-$  is correct

$\therefore$  Option (C)

65. Both can be distinguished by iodoform test.

$\text{Benzene ring}-COCH_3$  (Acetophenone) will perform the test


which req.  $NaOH + I_2 \rightarrow NaOI$   $\therefore$  Option (B)

66.  $pK_a \propto \frac{1}{\text{Acidic strength}}$

Acidic strength:  $HCOOH > Cl-CH_2CH_2COOH > CH_3CH_2COOH > CH_3CH_2OH$

$pK_a$  is inverse  $\therefore$  Option (A)

67.



It is soluble in Acid like HCl

but insoluble in base like NaOH  $\therefore$  Option (C)

68.

greater the basic strength, more

the reactivity towards HCl  $\therefore$  Option (C)

69.

Both "A" & "R" are correct and

"R" is correct explanation  $\therefore$  Option (A)

70.

Phenol is acidic so during azo

coupling its soln is prepared in NaOH  $\therefore$  Option (D)

71.

Conceptual  $\therefore$  Thymine is not present  $\therefore$  Option (B)

72.

73.

Due to deficiency of Vitamin C

i.e Ascorbic Acid causes scurvy

$\therefore$  Option (A)

74.

glucose  $\xrightarrow{NH_2OH}$  Its oxime

$\therefore$  Option (A)

75.

$$X_2 = 0.2 \therefore X_1 = 0.8$$

$$\frac{X_2}{X_1} = \frac{n_2}{n_1} = \frac{W_2 \times M_1 \times 1000}{M_2 \times W_1 \times 1000} = \frac{2}{8} = \frac{m \times 16}{1000}$$

$$\therefore m = 13.85$$


$\therefore$  ~~Convolving~~  $\therefore$   $\therefore$  Option (A)

76.

$A_{mix} > 0$  for ethanol + Acetone  $\therefore$  Option (C)

77.

n-octane is non-polar solvent so nonpolar solutes are soluble easily



$\therefore$  Option (77)

78. Hypertonic means higher concentration

0.9% w/v NaCl is isotonic

As  $1.2\% \frac{w}{v} > 0.9\% \frac{w}{v}$  ;  $1.2\% \frac{w}{v}$  will be hypertonic

----- Option (B)

79.

$$\Delta^{\circ}_{H_2O} = \lambda^{\circ}_{H^+} + \lambda^{\circ}_{OH^-}$$

$$= \Delta^{\circ}_{HCl} + \Delta^{\circ}_{NaOH} - \Delta^{\circ}_{NaCl}$$

$$= \lambda^{\circ}_{H^+} + \lambda^{\circ}_{Cl^-} + \lambda^{\circ}_{Na^+} + \lambda^{\circ}_{OH^-} - \lambda^{\circ}_{Na^+} - \lambda^{\circ}_{Cl^-}$$

-----  $\therefore$  Option (A)

80.

$$K_c = \text{A.L. of } \frac{n \times E^{\circ}_{\text{cell}}}{0.059}$$

$$= \frac{n \times E^{\circ}_{\text{cell}}}{0.059}$$

$$= 10$$

$$= \frac{2 \times 1.1}{0.059}$$

$$= 10$$

-----  $\therefore$  Option (D)